

# **Bioacoustic Absorption Spectroscopy**

Dr. Orest Diachok  
Acoustics Division, Code 7120  
Naval Research Laboratory  
Washington, D.C., 20375  
telephone: 202 404 4650    fax: 202 404 7813    e-mail: [orest@wave.nrl.navy.mil](mailto:orest@wave.nrl.navy.mil)

Document number: N0001400WX20881

## **LONG TERM GOAL**

Demonstrate the potential of bioacoustic absorption spectroscopy for tomographic mapping of the bioacoustic parameters of fish with swim bladders in shallow water.

## **OBJECTIVES**

Develop a propagation model that accounts for the effects of bioacoustic absorptivity on transmission loss in shallow water at frequencies between 0.2 and 10.0 kHz. Develop a bioacoustic model that accounts for the resonance frequencies of absorption lines, which are due to dispersed pelagic fish, and schools of pelagic fish with swim bladders. Develop a model of the seasonal variability of resonance frequencies. Demonstrate consistency between absorptivity and echo sounder based estimates of number densities.

## **APPROACH**

Design, construct and test an ultra broadband (0.2 – 10 kHz), light weight, long term, autonomous sources and receivers, that will permit long term monitoring of bioacoustic parameters. Conduct a series of bioacoustic absorptivity experiments in littoral seas in co-operation with fisheries biologists. These experiments will be designed to investigate the “signatures” of the two major classes of fish with swim bladders in time frequency space: physostomes (which have volumes that vary inversely with pressure) and physoclists (which have volumes that are independent of pressure).


Develop a transmission loss model that includes multiple bioacoustic absorbing layers with realistic shapes. Simulate effects of multiple absorbing layers on transmission loss vs. time and frequency of physostomes and physoclists. Apply model to new and previously published bioacoustic absorptivity measurements, derive bioacoustic parameters of sardines and anchovies and other fish, and demonstrate consistency with trawling data.

Invert bioacoustic parameters of anchovies from previously reported back-scattering data, which were made in concert with concurrent trawls in the seas off California, by matching theoretical computations with data. Demonstrate consistency between absorptivity and back-scattering based derivations of bioacoustic parameters. Continue to refine analysis of the causes of the differences between absorptivity and echo sounder based estimates of number densities of sardines in the Gulf of Lion.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2002</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2002 to 00-00-2002</b>	
4. TITLE AND SUBTITLE <b>Bioacoustic Absorption Spectroscopy</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Acoustics Division, Code 7120,,Naval Research Laboratory,,Washington,,DC, 20375</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <b>Demonstrate the potential of bioacoustic absorption spectroscopy for tomographic mapping of the bioacoustic parameters of fish with swim bladders in shallow water</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>6</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## WORK COMPLETED:

I served as Chief Scientist of the first bioacoustic absorption spectroscopy (BAS I) experiment in the Santa Barbara Channel. The experiment was conducted in concert with the NMFS Southwest Fisheries Science Center at the end of September. We recorded 3 days of high quality broadband acoustic data. The broadband source and the receiving arrays worked well. The source and the receiving arrays were easy to deploy and recover (deployment: ~ 60 minutes and recovery: ~ 30 minutes). The signal to noise ratio was high over most of the frequency range. The trawling was successful. The site was dominated by anchovies, as expected. Fish concentrations were initially low, increased with time, and were quite high on the third day of the experiment. SWFSC scientist, David Demer's, analysis of multi-frequency echo sounder data permitted isolation of fish from plankton, and through near real time analysis of target strengths provided an independent means for estimating fish lengths. Calculations of resonance frequencies for anchovy lengths derived from trawling and TS data are shown in Figure 1.




**Figure 1. Calculated resonance frequencies of anchovies, based on a preliminary analysis of trawling (13 and 9 cm) and TS (14, 10 and ~ 6 cm) measurements during BAS I. The minimum mesh size was 1.25 cm; the frequencies of echo sounders were 38 and 120 kHz.**

The peak at 13 cm corresponds to adult anchovies, and is consistent with historical measurements of this parameter. The peaks at 6 and 9 cm hypothetically correspond to “episodic” spawnings of juvenile anchovies, which were about 6 and 7 months old respectively (Butler, 1989). This is consistent with previously published bounds on lengths of juvenile anchovies in the Santa Barbara Channel in late September (5 – 10 cm).

## RESULTS

In preparation for BAS I I analyzed Van Holliday's backscattering measurements from dispersed anchovies. Matching of calculations and measurements resulted in estimates of the relative number densities and ensemble Q's of year classes. Results, which are shown in Figure 2, suggest that juveniles are characterized by multiple absorption lines.



**Figure 2. Measurements (●) due to Holliday (1972) and calculations of back-scattered energy vs. frequency. Resonance frequencies of 4.45 (blue), 1.95 (red), 1.45 (blue) and 1.10 (green) kHz correspond to 0.25, 2.25, 3.25 and 4.25 year old dispersed anchovies. Heavy solid line represents the sum of contributions from year classes. Absorption lines at 3.6 and 5.0 were not modeled, in the interest of clarity. Closely spaced peaks at 3.6, 4.45 and 5 kHz are hypothetically associated with juvenile anchovies.**

This interpretation is consistent with previously reported results, which indicate that spawning of anchovies is “episodic”. According to Butler (1989), several “episodes” generally occur in each spawning season. A similar pattern of absorption lines may be expected in BAS I data. A refereed paper that documents the results of this analysis will be published in the Journal of the Acoustical Society of America in 2001.

## **IMPACT / APPLICATIONS**

Naval significance: This research suggests that the detection range of naval tactical sonars may be significantly reduced when operating in shallow water environments where large numbers of pelagic fish are present. Strategically important areas, where fish concentrations may be particularly high, include the shallow seas off the coasts of the United States, Europe and China.

Fisheries applications: These results suggest that bioacoustic absorptivity can be used to estimate number density (biomass) of pelagic fish with swim bladders in littoral environments, and to classify fish by length.

## **TRANSITIONS**

Naval Research Laboratory (NRL) is supporting a three year 6.2 research program to conduct experimental and theoretical investigations on the effects of bioacoustic absorptivity on transmission loss in littoral seas. This research program started in FY 00.

## **RELATED PROJECTS**

Southwest Fisheries Science Center: biological sampling and fisheries sonar programs.

## **REFERENCES**

Holliday, V. Resonance structure in echoes from schooled pelagic fish, J. Acoust. Soc. Am., 51, 1322-1332, 1972.

Butler, J. 1989. Growth during the larval and juvenile stages of the northern anchovy, *Engraulis mordax*, in the California Current during 1980-84, Fishery Bulletin, U.S., 87, 645-652.

## **PUBLICATIONS**

Orest Diachok, Bernard Liorzou and Carla Scalabrin, "Estimation of the number density of fish from resonance absorptivity and echo sounder data", ICES Journal of Marine Science, 58, 137-153, 2001.

Orest Diachok, "Bioacoustic Absorption Spectroscopy: Estimation of the Biomass of Fish with Swim Bladders", Bioacoustics, 2001 (in press), and an *Invited* talk presented at the International Conference on Fish Bioacoustics, Chicago, Ill., 2001.

Orest Diachok and Stephen Wales, "Simulations of the potential of bioacoustic absorption spectroscopy for mapping length and depth distributions of fish in the ocean", in Proceedings of the International Conference on Acoustics, Rome, 2001.

Orest Diachok, "Interpretation of the spectra of energy scattered by dispersed anchovies", Journal of the Acoustical Society of America, 2001 (in press).

Orest Diachok, “Assessment of potential environmental effects for the bioacoustic absorption spectroscopy (BAS I) experiment”, Naval Research Laboratory document published on the Internet, 2001.

Orest Diachok, “Comparison of measured and calculated absorption spectra due to ensembles of fish with swim bladders” (Abstract), 140th Meeting of the Acoustical Society of America, 2000.

Orest Diachok and Stephen Wales, “Effects of Bioacoustic Absorptivity on Transmission Loss in the Yellow Sea” (Abstract) submitted to the US Navy Journal of Underwater Acoustics (*Invited*)

Orest Diachok, “Absorption spectroscopy: a new approach to estimation of biomass”, Fisheries Research, 47, 231-244, 2000; and *Invited talk*, presented at the 1997 Meeting of the American Fisheries Society.

Orest Diachok, “Bioacoustic absorptivity and transmission loss in littoral seas” (Abstract), 139th Meeting of the Acoustical Society of America, 2000.

Orest Diachok, “Effects of absorptivity due to fish on transmission loss in shallow water”, Journal of the Acoustical Society of America, 105, 2107-2128, 1999.

## **PATENTS**

The Naval Research Laboratory has applied for an international patent on my design of low cost, large bandwidth, lightweight, autonomous source and receiver arrays. These systems are specifically designed to permit bioacoustic absorption spectroscopy measurements to be made between a fixed broadband source and multiple, widely spaced fixed receiving arrays in littoral seas.